

ehp

ehponline.org

Environmental Health

P E R S P E C T I V E S

Published by the National Institute of
Environmental Health Sciences

Interaction of the Onset of Spring and Elevated
Atmospheric CO₂ on Ragweed (*Ambrosia
artemisiifolia* L.) Pollen Production

Christine A Rogers, Peter M Wayne, Eric A Macklin,
Michael L Muilenberg, Christopher J Wagner,
Paul R Epstein, and Fakhri A Bazzaz

doi:10.1289/ehp.8549 (available at <http://dx.doi.org/>)
Online 9 February 2006



The National Institute of Environmental Health Sciences
National Institutes of Health
U.S. Department of Health and Human Services

**Interaction of the Onset of Spring and Elevated Atmospheric CO₂ on
Ragweed (*Ambrosia artemisiifolia* L.) Pollen Production**

**Christine A Rogers¹, Peter M Wayne², Eric A Macklin³, Michael L Muilenberg¹,
Christopher J Wagner¹, Paul R Epstein⁴, Fakhri A Bazzaz⁵**

Address:

Harvard School of Public Health
Dept of Environmental Health
EER, Landmark Center PO15677
401 Park Drive, Boston, MA 02215

¹ Department of Environmental Health, Harvard School of Public Health, Boston, MA
crogers@hsph.harvard.edu, mmuil@hsph.harvard.edu

² New England School of Acupuncture, Watertown, MA pwayne@nesa.edu

³ New England Research Institutes, Watertown, MA, EMacklin@neri.org

⁴ Center for Health and the Global Environment, Harvard Medical School, Boston, MA
paul_epstein@hms.harvard.edu

⁵ Organismic and Evolutionary Biology, Harvard University, Cambridge, MA.

Author for Correspondence (by express mail):

Christine A. Rogers
Exposure Epidemiology and Risk Program
Harvard School of Public Health
Landmark Center, PO15677, Rm 428C
401 Park Drive, Boston, MA 02215

(617) 384-8814
(617) 384-8859 fax
crogers@hsph.harvard.edu

Running Title : Spring onset and CO₂ influence ragweed pollen

Keywords : Allergenic pollen, *Ambrosia artemisiifolia*, Climate change, Climate variability, Elevated CO₂, Global warming, Ragweed, Springtime warming

Article descriptor: Climate change

Acknowledgements

This study was funded by NOAA grant # NA06GP0673, through collaborative funding from NOAA, EPA, NASA, NSF and EPRI. Additional support was provided by a grant from the NIEHS, number P30 ES000002. The views expressed are solely those of the authors. The authors declare no competing financial interests.

Abbreviations:

CO₂ – carbon dioxide

NPK – nitrogen phosphorous potassium

ppm – parts per million

RCF – relative centrifugal force

rpm – revolutions per minute

Section Headers:

ABSTRACT

INTRODUCTION

MATERIALS AND METHODS

RESULTS

DISCUSSION

CONCLUSION

REFERENCES

TABLES

FIGURE LEGENDS

FIGURES

ABSTRACT

Increasing atmospheric CO₂ is responsible for climate changes that are having widespread effects on biological systems. One of the clearest changes is earlier onset of spring and lengthening of the growing season. We designed the current study to examine the interactive effects of timing of dormancy release of seeds with low and high atmospheric CO₂ on biomass, reproduction and phenology in ragweed plants (*Ambrosia artemisiifolia* L.), which produce highly allergenic pollen.

We released ragweed seeds from dormancy at three 15-day intervals and grew plants in climate-controlled glasshouses at either ambient or 700 ppm CO₂ concentrations, placing open-top bags over inflorescences to capture pollen. Measurements of plant height and weight, inflorescence number, weight, and length, and days to anthesis and anthesis date were made on each plant, and whole plant pollen productivity was estimated from an allometric-based model.

Timing and CO₂ interacted to influence pollen production. At ambient CO₂, the earlier cohort acquired a greater biomass, a higher average weight per inflorescence, a larger number of inflorescences, flowered earlier, and had 54.8% greater pollen production than the latest cohort. At high CO₂, plants showed greater biomass and reproductive effort compared to those in ambient CO₂ but only for later cohorts. In the early cohort, pollen production was similar under ambient and high CO₂, but in the middle and late cohorts, high CO₂ increased pollen production by 32% and 55% respectively compared to ambient CO₂ levels. Overall, ragweed pollen production can be expected to increase significantly under predicted future climate conditions.

INTRODUCTION

Global climate changes, driven by increased concentrations of greenhouse gases such as CO₂, are having widespread impacts on biotic systems including both direct and indirect effects on human health (Epstein 1999; Patz et al. 2005). One of the most dramatic effects of climate change seen thus far is on the timing of reproductive processes in plants (Fitter and Fitter 2002), including wind-pollinated types many of which have highly allergenic pollen (Clot 2003; van Vliet et al. 2002). Hence, predicted increases in CO₂, coupled with further changes in climate, could have important implications for individuals with allergies and asthma.

Many regions are currently experiencing warming effects associated with global climate change, including longer growing seasons and earlier arrival of spring (IPCC 2001; Karl and Trenberth 2003; Menzel 2000). These changes have already greatly impacted plant and animal populations by significantly influencing inter-annual population dynamics and phenology (Loeuille and Ghil 2004; Root et al. 2003). Analysis of temporal events since the 1950s, across a wide array of plant and animal species, indicates that spring phenology in northern temperate zones is advancing about 5 days each decade (Root et al. 2003). The trend toward earlier spring onset is particularly evident in early spring flowering of wind-pollinated tree species for which reproductive development and bud-burst in spring are highly temperature sensitive (Clot 2003; van Vliet et al. 2002). However, early spring onset may also affect temperature-dependent processes occurring over the entire growing season, not just those in early spring. For example, an early spring could also influence developmental and reproductive processes in later flowering plants.

While atmospheric CO₂ has no apparent direct effect on human health, it does have well known direct effects on plants. Plants grow larger, use water more effectively, and reach

maturity faster when grown in elevated CO₂ (Bazzaz 1990; Drake et al. 1997). In addition, a number of recent studies suggest that plants can also have an enhanced reproductive effort (Jablonski et al. 2002; LaDeau and Clark 2001; Stiling et al. 2004). These effects are generally thought to be beneficial in agriculture (Southworth et al. 2002); however, some studies suggest this enhanced reproductive effort can also lead to an increase in pollen production (Wayne et al. 2002; Ziska and Caulfield 2000). Therefore, global warming is a public health concern because it has the potential to alter the timing and abundance of aeroallergens (Beggs 2004), which could result in increased symptoms in those with allergic rhinitis or asthma.

An under-appreciated but important consideration is the interactive effects of CO₂ with other known or predicted changes in climate and their impact on biotic systems. For example, CO₂ may be driving the warming that results in earlier springs, but plants will experience both effects at the same time (i.e. higher CO₂ and a longer growing season). Hence, it is important to study how climate variables will interact to drive plant responses.

In this study we seek to increase our understanding of the potential response of common ragweed (*Ambrosia artemisiifolia* L.), a late-season flowering allergenic plant, to springtime climate variability, and examine interactive effects of increased CO₂. We performed a controlled environment study with simulated changes in the timing of spring, at both ambient and future predicted CO₂ levels, to test whether: 1) variability in the onset of spring alters the rate and magnitude of ragweed development, flowering phenology, and seasonal pollen production; and whether 2) atmospheric CO₂ concentrations directly alter ragweed development and productivity, and influence plant responses to climatic variability.

MATERIALS AND METHODS

Common ragweed (*A. artemisiifolia*) is a C₃ plant common to roadsides and disturbed habitats throughout most of the United States and Canada (Bassett and Crompton 1975). It is monoecious, with separate male and female flowers borne on the same plant on distinct axillary branches allowing for independent control of allocation to sexes (Payne 1963).

Seeds of *A. artemisiifolia* collected from wild populations in Woodstock, Illinois were vernalized by sowing seeds in six growth containers containing Pro-Mix compost (Red Hill, PA) and storing in a refrigerator at 4° C until ready for germination. Two trays at a time were transferred from cold conditions to the glasshouses at three 15-day intervals, creating three temporal cohorts that would simulate variability in the onset of the growing season and would include anticipated advances of spring several decades into the future. One tray from each cohort was placed in 380 ppm (ambient) and the other at 700 ppm (elevated) CO₂ concentration. From each pair of trays, seedlings were chosen that all germinated on the same day; the germination dates (23 May 2002, 7 Jun 2002, and 22 Jun 2002) were also at 15-day intervals. The middle cohort approximates the germination date of plants in the Boston area (personal observation).

Approximately 15 days after their germination, we transplanted 24 seedlings from each tray into 6-dry-quart capacity growth containers (22.23 cm diameter × 21.59 cm deep). Soil in each container was composed of a 4:1 mix of Pro-Mix compost (Red Hill, PA) and washed sand (Quickrete Co., Atlanta, GA). The soil mixture was amended with slow-release 14:14:14 NPK fertilizer (Osmocote, Scott's, Marysville, OH), and plants were watered daily.

The glasshouses are comprised of six modules structured as three blocks, each block having two modules of differing CO₂ concentrations (380 and 700 ppm). Containers were arranged in the modules according to their CO₂ and temporal cohort (i.e., 8 plants in each of

three temporal cohorts, in each of three glasshouse modules, at both low and high CO₂, for a total of 144 plants). Day/night temperatures were maintained at 26/21° C. Ambient glasshouse light levels were approximately 70% of full sun, supplemented with 6 hours of light daily (10 AM to 4 PM) from overhead metal halide lamps, thus allowing plants to experience natural variation in day length. Temperature, CO₂, and light were computer-controlled for all modules, and we used corn plants (*Zea mays*) to help maintain a constant CO₂ concentration in the low CO₂ modules. In each module, temporal cohorts of ragweed plants were separated, and the positions of the containers within each treatment were randomized at intervals to minimize edge effects. , Cohorts were grown at a foliar density of approximately 9 plants/m². We recorded measurements of flower phenology and date of first pollen release for each ragweed plant throughout the experiment.

We chose five male floral spikes at random from each plant in the first two cohorts and three from each plant in the third cohort at each CO₂ level and placed a 5 cm × 25 cm polyethylene bag over each selected spike, similar to the procedure used in Ziska and Caulfield (2000). On one side of the bag near the bottom, we cut a small slit and placed the spike inside. The slit was then taped shut and the bag left to collect pollen shed by the spike, with the tops of the bags left open for ventilation. After pollen production had stopped, we measured the length of the bagged flower spikes, cut each at the base, and stored the spike in the collection bag at – 20° C until ready for evaluation. Bags in which water accumulated due to watering or heavy condensation were discarded leaving 477 individual inflorescences.

After senescence, we harvested plants over a 3-day period from 16-18 September. Plant height and number of inflorescences were recorded, the plants were cut at the base, and all flower spikes removed and placed in bags separate from the vegetative material. We measured

the length of each floral spike on each plant. Roots were washed clean of dirt and also placed in separate bags. All plant material was dried at 70° C for 48 hours and we recorded separate dry weight measurements for all roots, flowers, and vegetative material.

For each bagged flower spike, pollen was recovered by twice repeated 30 second vortexing in a wash solution (distilled water with 0.02% Tween 20) in 15 ml Falcon tubes, followed by five minute centrifugation (2500 rpm; RCF=600). Pollen recoveries from the spike, and pollen rinsed from the polyethylene bag, were combined in a total volume of 2.0 ml wash solution. We determined the number of pollen grains per spike by calculating the pollen concentration in the wash suspension from microscopic counts using a glass hemacytometer (Hausser Scientific, Horsham, PA).

For each inflorescence, we estimated pollen production p_{ij} from an allometric model based on log inflorescence length, time of dormancy release, CO₂ concentration, total number of inflorescences, total weight of inflorescences, and days to anthesis,

$$[1] \quad \log(p_{ij}) = \mu + t_r + l_j + (tl)_{jr} + c_q + n_i + (cn)_{iq} + w_i + a_i + (al)_{ij}$$

where μ is a constant and j indexes each inflorescence of log length l_j on plant i with number of inflorescences n_i , total inflorescence weight w_i , and days to anthesis a_i , dormancy release at time t_r , and grown under CO₂ concentration c_q . Additional interaction terms did not improve model prediction. We estimated whole-plant pollen production p_i as the sum of pollen production over all inflorescences on each plant.

We used a two-way factorial design with time of dormancy release crossed with CO₂ treatment and CO₂ nested within greenhouse wing to assess the responses to the timing of dormancy release and CO₂ and we modeled estimated pollen count, inflorescence number, inflorescence weight, aboveground biomass, plant height, days to anthesis, and date of anthesis.

Time was included as a fixed term. Greenhouse wing and CO₂ within wing were included as random terms to permit broad inference. The Time × CO₂ interaction was included as a fixed term since plants were individually randomized.

RESULTS

The impact of variability in the onset of spring under scenarios of ambient and elevated CO₂ was assessed through several biomass (plant height, aboveground biomass), phenological (days to anthesis, anthesis date) and reproductive measures (number of inflorescences, inflorescence length, total weight of inflorescences, and pollen production).

The model estimating whole-plant pollen production explained 62% of the variation in measured pollen counts from 477 inflorescences collected from 141 of 144 individual plants. Pollen production per inflorescence was most strongly associated with inflorescence length, number of inflorescences per plant, and days to anthesis (Table 1). The negative association between inflorescence number and pollen production suggests a tradeoff with some plants producing fewer pollen-rich inflorescences and others producing more inflorescences each producing less pollen per unit length.

We examined the interaction of time of dormancy release and CO₂ concentration, and the results are presented in Table 2. Significant Time × CO₂ interaction terms were found in each of estimated pollen count, inflorescence number, inflorescence weight, aboveground biomass, and plant height (marginally). CO₂ treatment did not significantly affect days to anthesis or anthesis date.

We calculated least square means for each level of time and CO₂ along with 95% confidence limits for the measures of biomass, reproductive effort, phenology, and pollen productivity.

Plants in early spring cohorts had significantly greater aboveground biomass and height than the late cohort as shown in Figure 1, with the greatest difference between early and late cohorts. Little additional gain in biomass or height was achieved between middle and early cohorts under either CO₂ condition perhaps indicating that near maximal growth occurred in the longer growing seasons. However, in cohorts released from dormancy later, plants grown at elevated CO₂ acquired significantly greater height and weight. Hence, elevated CO₂ appears to have a greater impact on increasing biomass when plants are younger and/or smaller.

Earlier release from dormancy also increased reproductive effort measured by the number of inflorescences and inflorescence weight (Figure 2). There was a continuous trend toward a greater number of, and heavier, inflorescences in the middle and early cohorts at ambient CO₂ levels. In the early cohort, there was no difference in reproductive effort for plants grown at ambient versus high CO₂. Interestingly, at high CO₂, plants from the middle cohort had the highest number and heaviest inflorescences. At high CO₂, inflorescences were significantly heavier and more abundant in both the middle and later cohorts than at ambient CO₂.

The influence of growing season length and CO₂ on phenological responses of days to anthesis and anthesis date was also examined (Figure 3). Photoperiodic control of flower initiation is well documented in *A. artemisiifolia* and is similar to many other late-summer flowering plants. Therefore, little difference in the number of days to flowering (anthesis) and anthesis date was expected. Logically, plants released from dormancy earlier had a longer time until flowering. However, surprisingly, the anthesis date (date on which first pollen release was recorded) differed between the three cohorts. There was a consistent trend toward a later date of first anthesis in the later cohorts. Therefore, while flower initiation is reportedly under

photoperiodic control (Lewis et al. 1991), anthesis apparently is not. There was no effect of CO₂ on the number of days until anthesis or anthesis date for any cohort.

Finally, at ambient CO₂, estimates of whole plant pollen production, based on parameters outlined in the model above, were higher in earlier cohorts (Figure 4). At ambient CO₂, the simulated early spring cohort produced 54.8% more pollen compared to plants released from dormancy late. High CO₂ did not further increase pollen production relative to ambient CO₂ in the early cohort, but increased pollen production was observed in the middle (32.0% increase, $p = 0.0506$) and late (55.0% increase, $p = 0.0240$) cohorts.

DISCUSSION

This study is the first to assess the potential impact of earlier arrival of spring, and the interaction with CO₂, as expected with global warming and increased climate variability, on pollen productivity in allergenic plants. Based on the current rate of phenological advances (5 days/decade) (Root et al. 2003), the degree of advancement used in this study is similar to what might be expected 3 to 6 decades in the future. Our simulated effect of earlier spring dormancy release allowed ragweed plants to accumulate more resources through the season thereby increasing biomass and reproductive effort. Plants in ambient CO₂ released from dormancy earlier had increased height and weight, more and heavier inflorescences, and 54.8% higher pollen production compared to those released 30 days later. It should be noted that increased temperatures, which would accompany earlier spring and elevated CO₂ under future climate regimes, although not studied in these experiments, might also impact pollen production.

Because increasing atmospheric CO₂ is assured for the next several decades, and it is unknown how CO₂ might interact with climatic variables to influence plant responses, we also

determined the additional interactive effects of elevated CO₂ with variations in the onset of spring. We found that there was no additional advantage to plants in the earliest cohort grown under high CO₂. However, the number and weight of inflorescences was significantly greater at high CO₂ relative to ambient for plants in both the middle and late cohorts. Increased biomass and pollen production was also significantly higher in the late cohort at high CO₂. Hence, the reproductive disadvantage of a shorter growing season could be ameliorated when plants are grown at elevated CO₂.

It is a well known phenomenon of chamber studies that the advantage of elevated CO₂ is greatest early in plant development but diminishes over time (Drake et al. 1997). In essence, plants exhibit acclimation to elevated CO₂ with age (Long et al. 2004) or perhaps as a result of resource depletion due to the confines of growth within pots (Drake et al. 1997). We found the least difference in productivity between plants in ambient and elevated CO₂ for the earliest cohort. Early cohort plants at elevated CO₂ may have had an early advantage but then acclimated over time, and/or, the longer growing season may have been sufficient for ambient CO₂ plants to make up the early difference. In contrast, in the latest cohort with the shortest growing season, and hence the least amount of time for ambient CO₂ plants to make up the early difference in productivity, elevated CO₂ plants had significantly greater biomass, number and weight of inflorescences, and pollen production relative to ambient CO₂ plants. These results highlight the importance of examining the interactive effects of CO₂ with other climate variables in order to understand the implications of climate change.

The climate variability that stems from global warming is a significant concern. Our results show that variability in the onset of spring, elicits a strong increase in pollen production in early seasons at ambient CO₂ concentrations. However, at elevated CO₂, while pollen

productivity is enhanced, it is less sensitive to variability in season onset. Hence, in future climates with elevated CO₂, we predict pollen production will be just as robust in years with late springs as those with early springs. Overall, pollen production in ragweed can be expected to increase significantly under predicted future climate conditions.

Our results are consistent with the findings of other greenhouse and chamber studies on ragweed which have shown a 60-90% increase in pollen productivity with elevated CO₂ (700 or 600 ppm) compared to current ambient levels (Wayne et al. 2002; Ziska and Caulfield 2000). Of course, the ability to generalize results of closed environment experiments to natural field populations is an important issue; however, *A. artemisiifolia* also appears to be a strong competitor in mixed populations in elevated CO₂ (Bazzaz and Garbutt 1988). In addition, similar results on pollen productivity have been found in field studies. In cities, because of proximity to industrial and vehicular sources, atmospheric CO₂ concentrations and temperatures are much higher than the surrounding rural areas (Idso et al. 2002). Using a naturally occurring gradient in Baltimore where temperature and CO₂ are elevated by 1.8-2.0°C and 30% (to ~500ppm) respectively compared to outlying areas, Ziska et al. (2003) found that in experimental ragweed plots, plants increased biomass and pollen production with the degree of urbanization. This brings to light two facts; plants in the field are responding similarly to glasshouse experiments, and urban plants are currently experiencing changed atmospheric conditions that are likely altering their pollen productivity now, not decades into the future.

Both allergies and asthma have been increasing worldwide in recent decades, significantly above that expected from better diagnosing or increased reporting (von Mutius 1998). While the trend may be showing early signs of leveling off (Hertzen and Haahtela 2005; Lawson and Senthilselvan 2005), there still is a much greater proportion of the population that is

vulnerable to allergen exposure than ever before. Ragweed pollen allergens are some of the most potent in North America, and roughly 10% of the population is sensitized (Gergen et al. 1987). Diesel particles from truck and vehicle exhaust have been shown to act synergistically with pollen allergens in exacerbation of disease (Hauser et al. 2003) and are now thought to be an important factor in the recent rise in allergic disease (Riedl and Diaz-Sanchez 2005). Hence, an important question is whether greater ragweed pollen production (with or without diesel particle co-exposure) will lead to an increase in the frequency or severity of asthma and allergy symptoms, or to new sensitizations and a further increase in development of allergic disease.

CONCLUSION

The effects of global warming are complex, but studies of their impact on biotic communities clearly point toward secondary effects that could be detrimental to human health. Our study of *Ambrosia artemisiifolia* under conditions that simulate future levels of atmospheric CO₂ and increased temperatures shows that one effect—increased production of allergenic pollen—could strongly impact the significant proportion of the population with pollen allergies as climate change progresses. Since the results of this study suggest that under future conditions of global warming and elevated CO₂, pollen seasons will be more intense and could start earlier than expected, pollen forecasting and pollen avoidance strategies for sensitized individuals will be particularly important. Finally, we emphasize the importance of studying interactions between multiple predicted climate change parameters; in this case, the interactive effects of elevated CO₂ and variability in the onset of spring. Our study suggests that under future predicted greenhouse gas emissions and associated climate conditions, either an early spring onset, or variability in spring onset along with elevated CO₂, there will be an overall increase in ragweed pollen production.

REFERENCES

- Bassett IJ, Crompton CW. 1975. The biology of Canadian weeds. 11. *Ambrosia artemisiifolia* L. and *A. psilostachya* DC. Can J Plant Sci 55:463-476.
- Bazzaz FA. 1990. The response of natural ecosystems to the rising global CO₂ levels. Annu Rev Ecol Syst 21:167-196.
- Bazzaz FA, Garbutt K. 1988. The response of annuals in competitive neighborhoods: effects of elevated CO₂. Ecology 69:937-946.
- Beggs PJ. 2004. Impacts of climate change on aeroallergens: past and future. Clin Exp Allergy 34:1507-1513.
- Clot B. 2003. Trends in airborne pollen: An overview of 21 years of data in Neuchatel (Switzerland). Aerobiologia 19:227-234.
- Drake BG, Gonzalez-Meler MA, Long SP. 1997. More efficient plants: A consequence of rising atmospheric CO₂? Annu Rev Plant Physiol Plant Mol Bio 48:609-639.
- Epstein PR. 1999. Climate and health. Science 285:347-348.
- Fitter AH, Fitter RSR. 2002. Rapid changes in flowering time in British plants. Science 296:1689-1691.
- Gergen P, Turkeltaub P, Kovar M. 1987. The prevalence of allergic skin test reactivity to eight common aeroallergens in the US population: results from the second National Health and Nutrition Examination survey. J Allergy Clin Immunol 80:669-679.
- Hauser R, Rice TM, Krishna Murthy GG, Wand MP, Lewis D, Bledsoe T, et al. 2003. The upper airway response to pollen is enhanced by exposure to combustion particulates: a pilot human experimental challenge study. Environ Health Perspect 111:472-477.

- Hertzen L, Haahtela T. 2005. Signs of reversing trends in prevalence of asthma. *Allergy* 60:283-292.
- Idso SB, Idso CD, Balling Jr. RC. 2002. Seasonal and diurnal variations of near-surface atmospheric CO₂ concentration within a residential sector of the urban CO₂ dome of Phoenix, AZ, USA. *Atmos Environ* 36:1655-1660.
- IPCC. 2001. *Climate Change 2001: Synthesis Report*. Cambridge. A Contribution of Working Groups I, II, and III to the Third Assessment Report of the Intergovernmental Panel on Climate Change, Watson, R.T. and the Core Writing Team (eds.), Cambridge University Press, Cambridge UK and New York, NY, USA 398 pp.
- Jablonski LM, Wang X, Curtis PS. 2002. Plant reproduction under elevated CO₂ conditions: a meta-analysis of reports on 79 crop and wild species. *New Phytol* 156:9-26.
- Karl TR, Trenberth KE. 2003. Modern global climate change. *Science* 302:1719-1723.
- LaDeau SL, Clark JS. 2001. Rising CO₂ levels and the fecundity of forest trees. *Science* 292:95-98.
- Lawson JA, Senthilselvan A. 2005. Asthma epidemiology: has the crisis passed? *Curr Opin Pulm Med* 11:79-84.
- Lewis WH, Dixit AB, Wedner HJ. 1991. Asteraceae aeropollen of the western United States Gulf Coast. *Ann Allergy* 67:37-46.
- Loeuille N, Ghil M. 2004. Intrinsic and climatic factors in North-American animal population dynamics. *BMC Ecol* 4:6.
- Long SP, Ainsworth EA, Rogers A, Ort DR. 2004. Rising atmospheric carbon dioxide: plants FACE the future. *Annu Rev Plant Biol* 55:591-628.

- Menzel A. 2000. Trends in phenological phases in Europe between 1951 and 1996. *Int J Biometeorol* 44:76-81.
- Patz JA, Campbell-Lendrum D, Holloway T, Foley JA. 2005. Impact of regional climate change on human health. *Nature* 438:310-317.
- Payne WW. 1963. The morphology of the inflorescence of ragweeds (*Ambrosia-Franseria*: Compositae). *Am J Bot* 50:872-880.
- Riedl M, Diaz-Sanchez D. 2005. Biology of diesel exhaust effects on respiratory function. *J Allergy Clin Immunol* 115:221-228.
- Root TL, Price JT, Hall KR, Schneider SH, Rosenzweig C, Pounds JA. 2003. Fingerprints of global warming on wild animals and plants. *Nature* 421:57-60.
- Southworth J, Pfeifer R, Habeck M, Randolph J, Doering O, Rao D. 2002. Sensitivity of winter wheat yields in the midwestern United States to future changes in climate, climate variability, and CO₂ fertilization. *Clim Res* 22:73-86.
- Stiling P, Moon D, Hymus G, Drake B. 2004. Differential effects of elevated CO₂ on acorn density, weight, germination, and predation among three oak species in a scrub-oak forest. *Global Change Biol* 10:228-232.
- van Vliet A, Overeem A, de Groot R, Jacobs A, Spieksma FTM. 2002. The influence of temperature and climate change on the timing of pollen release in The Netherlands. *Int J Climatol* 22:1757-1767.
- von Mutius E. 1998. The rising trends in asthma and allergic disease. *Clin Exp Allergy* 28 Suppl 5:45-49.

Wayne P, Foster S, Connolly J, Bazzaz F, Epstein P. 2002. Production of allergenic pollen by ragweed (*Ambrosia artemisiifolia* L.) is increased in CO₂-enriched atmospheres. *Ann Allergy Asthma Immunol* 88:279-282.

Ziska LH, Caulfield FA. 2000. Rising CO₂ and pollen production of common ragweed (*Ambrosia artemisiifolia*), a known allergy-inducing species: implications for public health. *Aust J Plant Physiol* 27:893-898.

Ziska LH, Gebhard DE, Frenz DA, Faulkner S, Singer BD, Straka JG. 2003. Cities as harbingers of climate change: common ragweed, urbanization, and public health. *J Allergy Clin Immunol* 111:290-295.

Table 1. Regression coefficients for the model estimating whole-plant pollen productivity of *Ambrosia artemisiifolia*.

Parameter	Symbol	Level	Coeff	SE	t	p-value
Intercept	μ		3.47	1.63	2.13	0.0340
Log inflorescence length	l_j		5.48	0.652	8.40	<.0001
Time of dormancy release	t_r	Early	0.00	.	.	.
		Middle	1.63	0.370	4.41	<.0001
		Late	2.37	0.505	4.68	<.0001
Log inflorescence length \times Time of release	$(tl)_{jr}$	Early	0.00	.	.	.
		Middle	-0.665	0.151	-4.40	<.0001
		Late	-0.837	0.198	-4.24	<.0001
Number of inflorescences	n_i		-0.00503	0.000851	-5.91	<.0001
CO ₂ concentration	c_q	380 ppm	0.00	.	.	.
		700 ppm	-0.448	0.153	-2.93	0.0036
Number of inflorescences \times CO ₂	$(cn)_{iq}$	380 ppm	0.00	.	.	.
		700 ppm	0.00297	0.00106	2.79	0.0055
Weight of inflorescences	w_i		0.0509	0.0157	3.24	0.0013
Days to anthesis	a_i		0.119	0.0207	5.73	<.0001
Log inflorescence length \times Days to anthesis	$(al)_{ij}$		-0.0518	0.00840	-6.17	<.0001

Table 2. Effects of time of release, CO₂ concentration, and the interaction of Time and CO₂, were modeled on measures of biomass, reproduction, phenology, and pollen production.

Response	Term	F ^a	p-value.
Pollen count (estimated)	Time	8.49	0.0003
	CO ₂	2.54	0.2519
	Time × CO ₂	4.39	0.0143
Inflorescence number	Time	2.91	0.0579
	CO ₂	13.12	0.0685
	Time × CO ₂	3.58	0.0306
Inflorescence weight	Time	40.24	<.0001
	CO ₂	3.61	0.1979
	Time × CO ₂	8.66	0.0003
Aboveground biomass	Time	42.78	<.0001
	CO ₂	5.06	0.1534
	Time × CO ₂	4.13	0.0181
Plant height	Time	23.80	<.0001
	CO ₂	0.07	0.8125
	Time × CO ₂	2.97	0.0546
Days to anthesis	Time	62.40	<.0001
	CO ₂	1.63	0.3299
	Time × CO ₂	1.25	0.2890
Anthesis date	Time	49.42	<.0001

CO ₂	1.63	0.3299
Time × CO ₂	1.25	0.2890

a: for the F statistic, numerator degrees of freedom are Time=2, CO₂=1, Time × CO₂=2, and denominator degrees of freedom are Time=134, CO₂=2, Time × CO₂=134 (except for plant height where denominator df for Time and Time × CO₂ are 133).

FIGURE LEGENDS

Figure 1. End of season biomass measures of *Ambrosia artemisiifolia* a) aboveground biomass and b) plant height for three springtime dormancy release cohorts at 2 CO₂ concentrations (380 ppm and 700 ppm).


Figure 2. End of season reproductive measures of *Ambrosia artemisiifolia* a) number of inflorescences and b) inflorescence weight for three springtime dormancy release cohorts at 2 CO₂ concentrations (380 ppm and 700 ppm).

Figure 3. Phenology measures of *Ambrosia artemisiifolia* a) number of days to anthesis and b) anthesis date for three springtime dormancy release cohorts at 2 CO₂ concentrations (380 ppm and 700 ppm).

Figure 4. Pollen production in *Ambrosia artemisiifolia* for three springtime dormancy release cohorts grown at 2 CO₂ concentrations (380 ppm and 700 ppm).

Figure 1

a)



b)





Figure 2.

a)



b)





Figure 3.

a)



b)




Figure 4.

